

## 兔抗 AKT1(Ab-473)多克隆抗体

中文名称: 兔抗 AKT1(Ab-473)多克隆抗体

英文名称: Anti-AKT1(Ab-473) polyclonal antibody

别 名: AKT serine/threonine kinase 1; AKT; PKB; RAC; CWS6; PRKBA; PKB-ALPHA;

RAC-ALPHA

相关类别: 一抗

储 存: 冷冻(-20℃)

抗 原: AKT1(Ab-473)

宿 主: Rabbit

反应种属: Human, Mouse

标 记 物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格


**Background:** 

The serine-threonine protein kinase encoded by the A KT1 gene is catalytically inactive in serum-starved pri mary and immortalized fibroblasts. AKT1 and the relat ed AKT2 are activated by platelet-derived growth fact or. The activation is rapid and specific, and it is abrog ated by mutations in the pleckstrin homology domain of AKT1. It was shown that the activation occurs thro ugh phosphatidylinositol 3-kinase. In the developing n ervous system AKT is a critical mediator of growth fac tor-induced neuronal survival. Survival factors can sup press apoptosis in a transcription-independent manner by activating the serine/threonine kinase AKT1, which



## 全国订货电话 4008-723-722

	then phosphorylates and inactivates components of the apoptotic machinery. Mutations in this gene have been associated with the Proteus syndrome. Multiple alternatively spliced transcript variants have been found for this gene.
Applications:	WB, IHC
Name of antibody:	AKT1(Ab-473)
Immunogen:	Synthetic peptide of human AKT1
Full name:	AKT serine/threonine kinase 1
Synonyms:	AKT; PKB; RAC; CWS6; PRKBA; PKB-ALPHA; RAC-ALPH A
SwissProt:	P31749
IHC positive control:	Human breast carcinoma
IHC Recommend dilution:	50-100
WB Predicted band size:	56 kDa
WB Positive control:	293 and 3T3 cells
WB Recommended dilution:	500-1000





## 全国订货电话 4008-723-722

