

## 兔抗 BCR (Phospho-Tyr177)多克隆抗体

中文名称： 兔抗 BCR (Phospho-Tyr177)多克隆抗体

英文名称： Anti-BCR (Phospho-Tyr177) rabbit polyclonal antibody

别名： ALL; CML; PHL; BCR1; D22S11; D22S662

相关类别： 一抗

储存： 冷冻（-20℃） 避光

宿主： Rabbit

抗原： BCR (Phospho-Tyr177)

反应种属： Human, Mouse

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

#### Background:

A reciprocal translocation between chromosomes 2 and 9 produces the Philadelphia chromosome, which is often found in patients with chronic myelogenous leukemia. The chromosome 22 breakpoint for this translocation is located within the BCR gene. The translocation produces a fusion protein which is encoded by sequence from both BCR and ABL, the gene at the chromosome 9 breakpoint. Although the BCR-ABL fusion protein has been extensively studied, the function of the normal BCR gene product is not clear. The protein has serine/threonine kinase activity and is a GTPase-activating protein for p21rac. Two transcript variants encoding different is

	oforms have been found for this gene.
<b>Applications:</b>	WB, IHC
<b>Name of antibody:</b>	BCR (Phospho-Tyr177)
<b>Immunogen:</b>	Synthetic peptide of human BCR (Phospho-Tyr177)
<b>Full name:</b>	breakpoint cluster region (Phospho-Tyr177)
<b>Synonyms :</b>	ALL; CML; PHL; BCR1; D22S11; D22S662
<b>SwissProt:</b>	P11274
<b>IHC positive control:</b>	Human tonsil tumor
<b>IHC Recommend dilution:</b>	50-100
<b>WB Predicted band size:</b>	210 kDa
<b>WB Positive control:</b>	K562 cells untreated or treated with H2O2
<b>WB Recommended dilution:</b>	500-1000



