

## 兔抗 CCDC102B 多克隆抗体

中文名称：兔抗 CCDC102B 多克隆抗体

英文名称：Anti-CCDC102B rabbit polyclonal antibody

别名：coiled-coil domain containing 102B; AN; ACY1L; HsT1731; C18orf14

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：CCDC102B

反应种属：Human

标记物：Unconjugate

克隆类型：rabbit polyclonal

### 技术规格

<b>Background:</b>	CCDC102B (coiled-coil domain containing 102B), also known as AN, ACY1L or HsT1731, is a 513 amino acid protein that exists as three alternatively spliced isoforms. Widely expressed and found in multiple CNV (copy-number variant) regions, CCDC102B contains the deletion breakpoint of a maternally inherited deletion, which is 2.7?Mb in size, and maps to human chromosome 18q22.1. CCDC102B may play a role in the pathogenesis of diaphragmatic hernia, microphthalmia, colorectal carcinoma and schizophrenia. Encoding over 300 genes, chromosome 18 contains about 76 million bases. Translocation between chromosomes 18 and 14 is the most common translocation in cancers and occurs in follicular lymphomas. Niemann-Pick disease, here
--------------------	---

	editary hemorrhagic telangiectasia and erythropoietic protoporphyria are associated with chromosome 18.
<b>Applications:</b>	ELISA, WB
<b>Name of antibody:</b>	CCDC102B
<b>Immunogen:</b>	Full length fusion protein
<b>Full name:</b>	coiled-coil domain containing 102B
<b>Synonyms:</b>	AN; ACY1L; HsT1731; C18orf14
<b>SwissProt:</b>	Q68D86
<b>ELISA Recommended dilution:</b>	5000-10000
<b>WB Predicted band size:</b>	60 kDa
<b>WB Positive control:</b>	HEPG2, HUVEC and PC-3 cell lysates
<b>WB Recommended dilution:</b>	200-1000

