

## 兔抗 CDC40 多克隆抗体

中文名称: 兔抗 CDC40 多克隆抗体

英文名称: Anti-CDC40 rabbit polyclonal antibody

别名: EHB3; PRP17; PRPF17

储存: 冷冻 (-20℃) 避光

抗原: CDC40

宿主: Rabbit

反应种属: Human Mouse Rat

相关类别: 一抗

标记物: Unconjugate

克隆类型: Unconjugate

### 技术规格

<b>Background:</b>	Pre-mRNA splicing occurs in two sequential transesterification steps. The protein encoded by this gene is found to be essential for the catalytic step II in pre-mRNA splicing process. It is found in the spliceosome, and contains seven WD repeats, which function in protein-protein interactions. This protein has a sequence similarity to yeast Prp17 protein, which functions in two different cellular processes: pre-mRNA splicing and cell cycle progression. It suggests that this protein may play a role in cell cycle progression.
<b>Applications:</b>	WB
<b>Name of antibody:</b>	CDC40
<b>Immunogen:</b>	Synthesized peptide derived from internal of human CDC

	40.
<b>Full name:</b>	cell division cycle 40
<b>Synonyms :</b>	EHB3; PRP17; PRPF17
<b>SwissProt:</b>	O60508
<b>WB Predicted band size:</b>	66 kDa
<b>WB Positive control:</b>	K562 cells, Huvec cells and COLO cells lysates
<b>WB Recommended dilution:</b>	500-3000

