

兔抗 HLA-DPA1 多克隆抗体

中文名称: 兔抗 HLA-DPA1 多克隆抗体

英文名称: Anti-HLA-DPA1 rabbit polyclonal antibody

别名: major histocompatibility complex, class II, DP alpha 1; PLT1; HLADP; HLASB; DP(W3); DP(W4); HLA-DP1A

抗原: HLA-DPA1

储存: 冷冻 (-20°C)

宿主: Rabbit

反应种属: Human

相关类别: 一抗

标记物: Unconjugate

克隆类型: rabbit polyclonal

技术规格

Background:

HLA-DPA1 belongs to the HLA class II alpha chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DP A) and a beta (DPB) chain, both anchored in the membrane. It plays a central role in the immune system by presenting peptides derived from extracellular proteins. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The alpha chain is approximately 33-35 kDa and its gene contains 5 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and the cytoplasmic tail. Within the DP molecule both the alpha chain and the beta chain contain the polymorphisms specifying the peptide binding specificities.

	es, resulting in up to 4 different molecules.
Applications:	ELISA, WB, IHC
Name of antibody:	HLA-DPA1
Immunogen:	Synthetic peptide of human HLA-DPA1
Full name:	major histocompatibility complex, class II, DP alpha 1
Synonyms:	PLT1; HLADP; HLASB; DP(W3); DP(W4); HLA-DP1A
SwissProt:	P20036
ELISA Recommended dilution:	5000-10000
IHC positive control:	Human tonsil
IHC Recommend dilution:	40-200
WB Predicted band size:	29 kDa
WB Positive control:	Raji cell, Human spleen tissue lysates
WB Recommended dilution:	500-2000



