

# 兔抗 IKBKG(Ab-85) 多克隆抗体

中文名称： 兔抗 IKBKG(Ab-85) 多克隆抗体

英文名称： Anti-IBKKG(Ab-85) rabbit polyclonal antibody

别 名： IP; IP1; IP2; FIP3; IPD2; NEMO; FIP-3; Fip3p; IMD33; AMCBX1; ZC2HC9; IKK-gamma

相关类别： 一抗

储 存： 冷冻 (-20°C) 避光

宿 主： Rabbit

抗 原： IKBKG(Ab-85)

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

## 技术规格

<b>Background:</b>	This gene encodes the regulatory subunit of the inhibitor of kappaB kinase (IKK) complex, which activates NF-kappaB resulting in activation of genes involved in inflammation, immunity, cell survival, and other pathways. Mutations in this gene result in incontinentia pigmenti, hypohidrotic ectodermal dysplasia, and several other types of immunodeficiencies. Multiple transcript variants encoding different isoforms have been found for this gene.
<b>Applications:</b>	WB, IF
<b>Name of antibody:</b>	IBKKG(Ab-85)
<b>Immunogen:</b>	Synthesized non-phosphopeptide derived from human IKK-γ around the phosphorylation site of serine 85 (Q-A-S(p)-Q-R).
<b>Full name:</b>	inhibitor of kappa light polypeptide gene enhancer in B-cells

	, kinase gamma
<b>Synonyms :</b>	IP; IP1; IP2; FIP3; IPD2; NEMO; FIP-3; Fip3p; IMD33; AMCBX1; ZC2HC9; IKK-gamma
<b>SwissProt:</b>	Q9Y6K9
<b>WB Predicted band size:</b>	48 kDa
<b>WB Positive control:</b>	HepG2 cells treated with Anisomycin (0.5uM, 5hours) lysate
<b>WB Recommended dilution :</b>	500-3000
<b>IF positive control:</b>	HeLa cells
<b>IF Recommend dilution:</b>	100-500



