

CCL13 抗原（重组蛋白）

中文名称：CCL13 抗原（重组蛋白）

英文名称：CCL13 Antigen (Recombinant Protein)

储 存： 冷冻（-20℃）

相关类别： 抗原

概述

Fusion protein corresponding to a region derived from 17-98 amino acids of human CCL13

技术规格

Full name:	C-C motif chemokine ligand 13
Synonyms:	NCC1; CKb10; MCP-4; NCC-1; SCYL1; SCYA13
Swissprot:	Q99616
Gene Accession:	BC008621
Purity:	>85%, as determined by Coomassie blue stained SDS-PAGE
Expression system:	Escherichia coli
Tags:	His tag C-Terminus, GST tag N-Terminus
Background:	This antimicrobial gene is one of several Cys-Cys (CC) cytokine genes clustered on the q-arm of chromosome 17. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjacent cysteines. The cytokine encoded by this gene displays chemotactic activity for monocytes, lymphocytes, basophils and eosinophils, but not neutrophils. This chemokine plays a role in accumulation of leukocytes during inflammation. It may also be involved in the recruitment of monocytes into the arterial wall during atherosclerosis.