

## CCNA1 抗原（重组蛋白）

中文名称：CCNA1 抗原（重组蛋白）

英文名称：CCNA1 Antigen (Recombinant Protein)

储存：冷冻（-20℃）

相关类别：抗原

概述

Fusion protein corresponding to a region derived from 166-465 amino acids of human CCNA1

技术规格

<b>Full name:</b>	cyclin A1
<b>Synonyms:</b>	CT146
<b>Swissprot:</b>	P78396
<b>Gene Accession:</b>	BC036346
<b>Purity:</b>	>85%, as determined by Coomassie blue stained SDS-PAGE
<b>Expression system:</b>	Escherichia coli
<b>Tags:</b>	His tag C-Terminus, GST tag N-Terminus
<b>Background:</b>	The protein encoded by this gene belongs to the highly conserved cyclin family, whose members are characterized by a dramatic periodicity in protein abundance through the cell cycle. Cyclins function as regulators of CDK kinases. Different cyclins exhibit distinct expression and degradation patterns which contribute to the temporal coordination of each mitotic event. The cyclin encoded by this gene was shown to be expressed in testis and brain, as well as in several leukemic cell lines, and is thought to primarily function in the control of the germline meiotic cell cycle. This cyclin binds both CDK2 and CDC2 kinases, which give two distinct kinase activities, one appearing in S phase, the other in G2, and thus regulate separat

e functions in cell cycle. This cyclin was found to bind to important cell cycle regulators, such as Rb family proteins, transcription factor E2F-1, and the p21 family proteins. Multiple transcript variants encoding different isoforms have been found for this gene.