

## 兔抗 KLRC3 多克隆抗体

中文名称：兔抗 KLRC3 多克隆抗体

英文名称：Anti-KLRC3 rabbit polyclonal antibody

别名：NKG2E; NKG2-E

相关类别：一抗

储存：冷冻（-20℃）

宿主：Rabbit

抗原：KLRC3

反应种属：Human

标记物：Unconjugate

克隆类型：Unconjugate

### 技术规格

<b>Background:</b>	Natural killer (NK) cells are lymphocytes that can mediate lysis of certain tumor cells and virus-infected cells without previous activation. They can also regulate specific humoral and cell-mediated immunity. NK cells preferentially express several calcium-dependent (C-type) lectins, which have been implicated in the regulation of NK cell function. KLR C3 is a member of the NKG2 group which are expressed primarily in natural killer (NK) cells and encodes a family of transmembrane proteins characterized by a type II membrane orientation (extracellular C terminus) and the presence of a C-type lectin domain. The NKG2 gene family is located within the NK complex, a region that contains sev
--------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

	eral C-type lectin genes preferentially expressed on NK cells. Alternative splicing results in multiple transcript variants encoding different isoforms.
<b>Applications:</b>	ELISA, IHC
<b>Name of antibody:</b>	KLRC3
<b>Immunogen:</b>	Synthetic peptide of human KLRC3
<b>Full name:</b>	killer cell lectin-like receptor subfamily C, member 3
<b>Synonyms :</b>	NKG2E; NKG2-E
<b>SwissProt:</b>	Q07444
<b>ELISA Recommended dilution:</b>	5000-10000
<b>IHC positive control:</b>	Human tonsil and human cervical cancer
<b>IHC Recommend dilution:</b>	25-100



