

## 兔抗 CST2 多克隆抗体

中文名称：兔抗 CST2 多克隆抗体

英文名称： Anti-CST2 rabbit polyclonal antibody

相关类别： 一抗

抗 原： CST2

储 存： 冷冻（-20℃）

宿 主： Rabbit

反应种属： Human

标记物： Unconjugate

克隆类型： rabbit polyclonal

### 技术规格

<b>Background:</b>	<p>The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The type 2 cystatin proteins are a class of cysteine proteinase inhibitors found in a variety of human fluids and secretions, where they appear to provide protective functions. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. This gene is located in the cystatin locus and encodes a secreted thiol protease inhibitor found at high levels in saliva, tears and seminal plasma.</p>
<b>Applications:</b>	ELISA, WB, IHC

<b>Name of antibody:</b>	CST2
<b>Immunogen:</b>	Fusion protein of human CST2
<b>Full name:</b>	cystatin SA
<b>SwissProt:</b>	P09228
<b>ELISA Recommended dilution:</b>	2000-5000
<b>IHC positive control:</b>	Human cervical cancer and human thyroid cancer
<b>IHC Recommend dilution:</b>	50-200
<b>WB Predicted band size:</b>	16 kDa
<b>WB Positive control:</b>	Lovo cells
<b>WB Recommended dilution:</b>	200-1000



