

## 兔抗 SLC16A2 多克隆抗体

中文名称: 兔抗 SLC16A2 多克隆抗体

英文名称: Anti-SLC16A2 rabbit polyclonal antibody

别 名: AHDS; MCT7; MCT8; XPCT; MCT 7; MCT 8; MRX22; DXS128; DXS128E

抗 原: SLC16A2

储 存: 冷冻(-20℃) 避光

宿 主: Rabbit

相关类别: 一抗

反应种属: Human

标记物: Unconjugate

克隆类型: Unconjugate

## 技术规格

| Background:       | This gene encodes an integral membrane protein that functions as a transporter of thyroid hormone. The encoded protein facilitates the cellular importation of thyroxine (T4), triiodothyronine (T3), reverse triiodothyronine (rT3) and diidothyronine (T2). This gene is expressed in many tissues and likely plays an important role in the development of the central nervous system. Loss of function mutations in this gene are associated with psychomotor retardation in males while females exhibit no neurological defects and more moderate thyroid-deficient phenotypes. |
|-------------------|--|
| Applications:     | WB   |
| Name of antibody: | SLC16A2  |
| Immunogen:        | Synthesized peptide derived from internal of human SLC1 6A2.   |



## 全国订货电话 4008-723-722

| Full name:               | solute carrier family 16, member 2 (thyroid hormone trans porter) |
|--------------------------|---|
| Synonyms :               | AHDS; MCT7; MCT8; XPCT; MCT 7; MCT 8; MRX22; DXS12<br>8; DXS128E  |
| SwissProt:               | P36021  |
| WB Predicted band size:  | 60 kDa  |
| WB Positive control:     | K562 cells lysate   |
| WB Recommended dilution: | 500-3000  |

