

## 兔抗 STK40 多克隆抗体

中文名称: 兔抗 STK40 多克隆抗体

英文名称: Anti-STK40 rabbit polyclonal antibody

别 名: SHIK; SgK495

相关类别: 一抗

抗 原: STK40

储 存: 冷冻(-20℃)

宿 主: Rabbit

反应种属: Human, Mouse, Rat

标记物: Unconjugate

克隆类型: rabbit polyclonal

## 技术规格

	STK40 (Serine/threonine-protein kinase 40), also known
	as Sugen kinase 495, is a 435 amino acid protein kinas
	e that may be a negative regulator of NFкВ and p53-m
	ediated gene expression. Localized to both the cytoplas
Background:	m and the nucleus, STK40 is ubiquitously expressed. ST
	K40 has a central serine/threonine protein kinase domai
	n and is homologous to TRB-3, a protein that regulates
	activation of MAP kinases and inhibits NFkB-mediated
	gene transcription.
Applications:	ELISA, WB, IHC
Name of antibody:	STK40



Immunogen:	Fusion protein of human STK40
Full name:	serine/threonine kinase 40
Synonyms:	SHIK; SgK495
SwissProt:	Q8N2I9
<b>ELISA Recommended dilution:</b>	1000-2000
IHC positive control:	Human cervical cancer and human thyroid cancer
IHC Recommend dilution:	50-200
WB Predicted band size:	49 kDa
WB Positive control:	Human placenta tissue
WB Recommended dilution:	200-1000





