

兔抗 CD274 多克隆抗体

- 中文名称：兔抗 CD274 多克隆抗体
- 英文名称：Anti-CD274 rabbit polyclonal antibody
- 抗 原：CD274
- 别 名：B7-H; B7H1; PDL1; PD-L1; PDCD1L1; PDCD1LG1
- 储 存：冷冻（-20℃）
- 宿 主：Rabbit
- 相关类别：一抗
- 反应种属：Human, Mouse
- 标 记 物：Unconjugate
- 克隆类型：rabbit polyclonal

技术规格

Background:

This gene encodes an immune inhibitory receptor ligand that is expressed by hematopoietic and non-hematopoietic cells, such as T cells and B cells and various types of tumor cells. The encoded protein is a type I transmembrane protein that has immunoglobulin V-like and C-like domains. Interaction of this ligand with its receptor inhibits T-cell activation and cytokine production. During infection or inflammation of normal tissue, this interaction is important for preventing autoimmunity by maintaining homeostasis of the immune response. In tumor microenvironments, this interaction provides an immune escape for tumor cells through cytotoxic T-cell inactivation. Expression of this gene in tumor cells is considered to be prognostic in many types of human malignancies, including colon cancer and renal cell carcinoma. Alternative splicing results in

| | |
|------------------------------------|--|
| | multiple transcript variants. |
| Applications: | ELISA, WB |
| Name of antibody: | CD274 |
| Immunogen: | Synthetic peptide of human CD274 |
| Full name: | CD274 molecule |
| Synonyms: | B7-H; B7H1; PDL1; PD-L1; hPD-L1; PDCD1L1; PDCD1LG1 |
| SwissProt: | Q9NZQ7 |
| ELISA Recommended dilution: | 5000-10000 |
| WB Predicted band size: | 33 kDa |
| WB Positive control: | Human placenta tissue lysate |
| WB Recommended dilution : | 500-2000 |

kDa

130—
95—
72—
55—
36—
28—
17—

